Fluid Phase Equilibria, 34 (1987) 313–314
Elsevier Science Publishers B.V., Amsterdam – Printed in The Netherlands

Contents Volume 34 (1987)

Volume 34, No. 1

A three-parameter cubic equation of state for asymmetric mixture density calculations
J.-M. Yu and B.C.-Y. Lu (Ottawa, Ont., Canada) 1–19
A simple modification of the Flory–Huggins theory for polymers in non-polar or slightly polar solvents
A. Vetere (Milan, Italy) 21–35
Solubility measurements of naphthol isomers in supercritical CO_2 by a recycle technique
C.-S. Tan and J.-Y. Weng (Taiwan, Republic of China) 37–47
Excess enthalpy, excess heat capacity and excess volume of 1,2,4-trimethylbenzene+, and 1-methylnaphthalene+an n-alkane
E. Wilhelm (Vienna, Austria), A. Inglese (Bari, Italy), A.H. Roux and J.-P.E. Grolier (Aubière, France) 49–67
Solubility of propane and butane in mixtures of n-alkanes
R. González, F. Murrieta Guevara, O. Parra and A. Trejo (Mexico, Mexico) . 69–81
Vapor pressures of binary mixtures of carbon dioxide with benzene, n-hexane and cyclohexane up to 7 MPa
G.-I. Kaminishi, C. Yokoyama and S. Takahashi (Sendai, Japan) 83–100

Short Communication

Fluid phase stability and equations of state
I.F. Radzyminski and W.B. Whiting (Morgantown, WV, U.S.A.) 101–110
Book Reviews 111–112

Volume 34, Nos. 2–3

The generalized van der Waals partition function—IV. Local composition models for mixtures of unequal-size molecules
K.-H. Lee and S.I. Sandler (Newark, DE, U.S.A.) 113–147
Flash calculations for polydisperse fluids: a variational approach
A.G. Schlijper (Houston, TX, U.S.A.) 149–169
Perturbed hard sphere equations of state of real fluids—III. Residual parameter a_p of non-polar liquids
I. Nezbeda and K. Aim (Prague, Czechoslovakia) 171–188
An analysis of the industrial use of a phase equilibria prediction model based on thermodynamic perturbation theory
B. Moser and H. Kistenmacher (Höllriegelskreuth, F.R.G.) 189–201
Effect of cubic equations of state on enthalpy departure calculations
Y. Adachi and H. Sugie (Nagoya, Japan) 203–218

Application of the Kirkwood–Buff theory of solutions to dilute supercritical mixtures
H.D. Cochran (Oak Ridge, TN, U.S.A.), L.L. Lee and D.M. Pfund (Norman, OK, U.S.A.) 219–234

Thermodynamic properties of pure fluids and solutions from the hole group-contribution model
N.A. Smirnova and A.I. Victorov (Leningrad, U.S.S.R.) 235–263

Excess volumes of trichloroethylene with methylethylketone, diethylketone, methylisobutylketone, cyclohexanone and 1,4-dioxane at 298.15, 308.15 and 318.15 K
K. Purnachandra Rao and K.S. Reddy (Tirupati, India) 265–272

Fluid-phase equilibria of binary and ternary mixtures of 1-dodecanol with CO_2, $CClF_3$, CHF_3 and N_2 at 392 K and at pressures up to 90 MPa
G. Katzenski-Ohling and G.M. Schneider (Bochum, F.R.G.) 273–286

Measurements and calculations of phase equilibria of binary mixtures of ethane + eicosane. Part I: vapour + liquid equilibria
C.J. Peters, J.L. de Roo (Delft, The Netherlands) and R.N. Lichtenthaler (Heidelberg, F.R.G.) 287–308

Letter to the Editor

Density-dependent mixing rules for the calculation of fluid phase equilibria at high pressures
U.K. Deiters (Bochum, F.R.G.) 309–310

Author Index 311

Contents of Vol. 34 (1987) 313–314